

CD27 is required for generation and long-term maintenance of T cell immunity

Jenny Hendriks¹, Loes A. Gravesteyn¹, Kiki Tesselaar³, René A. W. van Lier³, Ton N. M. Schumacher² and Jannie Borst¹

The Traf-linked tumor necrosis factor receptor family member CD27 is known as a T cell costimulatory molecule. We generated CD27^{-/-} mice and found that CD27 makes essential contributions to mature CD4⁺ and CD8⁺ T cell function: CD27 supported antigen-specific expansion (but not effector cell maturation) of naïve T cells, independent of the cell cycle-promoting activities of CD28 and interleukin 2. Primary CD4⁺ and CD8⁺ T cell responses to influenza virus were impaired in CD27^{-/-} mice. Effects of deleting the gene encoding CD27 were most profound on T cell memory, reflected by delayed response kinetics and reduction of CD8⁺ virus-specific T cell numbers to the level seen in the primary response. This demonstrates the requirement for a costimulatory receptor in the generation of T cell memory.

The tumor necrosis factor (TNF) receptor family contains death receptors, as well as receptors that bind TNF receptor-associated factors (Trafs)¹. Many TNF receptor family members and their TNF-related transmembrane ligands are expressed on cells of the immune system where they play important roles. The Traf-linked receptor subgroup includes CD27 (also known as TNFRSF7), CD30, OX-40 (also known as CD134) and 4-1BB (also known as CD137), which selectively regulate lymphocyte function². Other prominent members are RANK and CD40, which additionally modulate the response of dendritic cells (DCs)^{3,4}. Traf-linked receptors have been implicated in cell proliferation, differentiation, survival and migration. For some receptors such as CD30, a death-inducing capacity has been observed, which may be conveyed by membrane TNF⁵. Trafs signal to NF-κB and Jun kinase, but the array of genes they target is undefined. It is proposed that Trafs counteract apoptosis⁶⁻⁸ via the inhibitor of apoptosis proteins (IAPs)⁹ and/or NF-κB¹⁰.

The relevance of Traf-linked receptors was initially ascertained by analyzing their expression patterns, as well as their functional effects *in vitro*. CD27, CD30, 4-1BB and OX-40 are exclusively expressed on cells of the lymphoid lineage, mostly in an activation-specific manner, and all enhance T cell receptor (TCR)-induced T cell expansion. CD27 and its ligand, CD70, have been defined at the protein, cDNA and genomic level in both human and mouse¹¹⁻¹⁶. CD27 is found on NK, T and B cell populations¹⁷. In humans and mice, the majority of CD4⁺ and CD8⁺ naïve peripheral T cells express CD27 and expression is up-regulated upon TCR stimulation¹⁸⁻²⁰. Loss of CD27 expression is irreversible and seems to represent terminal effector T cell differentiation^{19,21}. In humans, naïve B cells lack CD27 but antigen receptor stimulation induces expression. CD27⁺ B cells display all the functional and phenotypic char-

acteristics of memory cells^{22,23}. Whether the same is true in mice remains to be shown. Expression of the ligands for CD27 and related receptors is very restricted and tightly regulated by antigen and cytokines^{2,17,24,25}. The CD27 ligand CD70 is transiently up-regulated by antigen receptor stimulation on both T and B cells and thus reflects recent antigenic priming^{16,17}. CD70 has also been found on thymic DCs in humans²⁶ and on lymph node DCs from infected mice²⁷, but not on any other cell types.

CD27 ligation enhances TCR-induced expansion of both CD4⁺ and CD8⁺ peripheral T cells, but the nature of this costimulatory signal is ill-defined^{13,14,20,28,29}. One study documents that CD27-CD70 interaction promotes differentiation of CD8⁺ T cells to effector cytotoxic T cells (CTLs)³⁰. On B cells, CD27 ligation only marginally enhances expansion, but promotes plasma cell differentiation and immunoglobulin production³¹. The CD27-CD70 pair may contribute to T cell expansion through T-T cell, and possibly T cell-DC, interactions and to B cell differentiation through T-B cell and/or B-B cell interactions. Targeted deletion of receptors and ligands in mice is the optimal approach to determine the exact contribution each Traf-linked TNF receptor family member makes to the immune system. We present here the effects of targeting of the gene encoding CD27 (*Tnfrsf7*) on T cell development and peripheral responsiveness. Our analysis revealed that CD27 is required for effective generation and long-term maintenance of antigen-specific CD4⁺ and CD8⁺ T cell immunity.

Results

Generation of CD27-deficient mice

Genomic clone λ5, containing the complete coding region of *Tnfrsf7*, was isolated and characterized by Southern blotting and nucleotide sequencing (Fig. 1a). This led to the definite localization

Divisions of ¹Cellular Biochemistry and ²Immunology, The Netherlands Cancer Institute, ³Department of Immunobiology, Laboratory of the Netherlands Red Cross Blood Transfusion Service, 1066 CX Amsterdam, The Netherlands. Correspondence should be addressed to J. B. (jborst@nki.nl).




Figure 1. *Tnfrsf7*-targeting strategy and result. (a) Restriction map of the $\lambda 5$ clone encoding murine *Tnfrsf7* and the targeting construct derived from it. The dashed line indicates genomic DNA located outside of $\lambda 5$. Restriction sites indicated are: A: *Apa1*, B: *BamHI*, Bg: *BglII*, H: *HindIII*, K: *KpnI*, RV: *EcoRV*, S: *SphI*, Scal, Sm: *SmaI* and X: *XbaI*. Exons are depicted as wide filled bars (I–VI). The open reading frame of mCD27 starts between *BglII* and *SphI* sites and ends at the asterisk (*). (b) Southern blot of mouse-tail DNA cut with *Scal* and probed with the 3' probe indicated in a, visualizing a 9-kb WT *Tnfrsf7* fragment and a 7-kb mutated *Tnfrsf7* fragment. (c) Flow cytometric analysis of spleen cells from CD27^{+/+} mice and CD27^{-/-} mice. Cells were double stained as indicated.

lymphoid organs was normal in CD27^{-/-} mice. In the thymus, the size of CD4⁺CD8⁻, CD4⁺CD8⁺ and mature CD4⁺CD8⁻ and CD4⁺CD8⁺ populations, as well as subset composition of the CD4⁺CD8⁻ compartment were similar to WT (data not shown). Thus, CD27 was not required for antigen-independent expansion and differentiation of (precursor) T and B cells or for long-term survival of naïve lymphocytes.

Studies have implicated CD27 in costimulation of peripheral CD4⁺ and CD8⁺ T cells: TCR-induced [³H]thymidine incorporation is enhanced by monoclonal antibody (mAb) to CD27 or recombinant ligand^{13,14,20,28,29}. Conversely, mAbs to CD70 inhibit this response, indicating that CD27-CD70 interaction supports TCR-induced T cell expansion^{16,29}. Purified, naïve peripheral T cells from CD27^{-/-} mice showed significantly decreased TCR-induced [³H]thymidine incorporation as compared to WT T cells: 1.8-fold over a concentration range of mAb to CD3 ϵ (Fig. 2a). The same result was obtained when incorporation of radioactivity was assayed after 1 or 2 days of culture (data not shown). Recombinant interleukin 2 (IL-2) or costimulation with a mAb to CD28 increased the expansion of both WT and CD27^{-/-} cells, but did not rescue the defect in CD27^{-/-} cells (Fig. 2a). Thus, reduced [³H]thymidine incorporation was not caused by deficient IL-2 production and the CD27 costimulatory signal was qualitatively different from the CD28 signal.

The cell-permeant fluorescent dye carboxyfluorescein diacetate succinimidyl ester (CFSE) was used to study the effect of CD27 on cell cycle activity. Upon cell division, CFSE is equally distributed between daughter cells, allowing resolution of up to eight cycles of cell division by flow cytometry. Costimulation *via* CD28³² increased the proportion of T cells entering into cell cycle on TCR stimulation, as well as cell cycle activity (number of cycles completed within a 3-day time period) (Fig. 2b). However, absence of CD27 did not affect either cell cycle entry or activity (Fig. 2b). Similarly, costimulation of CD3-activated WT T cells with mAb to CD27 did

of exons I, II, V and VI. Exons V and VI are separated by an intron of 30 nucleotides and contain the end of the coding region and the 3' untranslated region. Exons III and IV were not localized exactly but are contained in the indicated restriction fragments. In the targeting construct, the complete *Tnfrsf7* coding region was replaced by the hygromycin-resistance gene. After selection, DNA from transfected embryonic stem (ES) cell clones was cut with *Scal* and hybridized with the 3' probe (Fig. 1a). Homologous recombination yielded a hybridizing fragment of 7 kb, whereas the wild-type (WT) fragment is 9 kb. A homologous recombinant ES cell clone was injected into C57BL/6 blastocysts. Resulting chimeric mice were crossed with FVB mice to attain germ line transmission of the mutated *Tnfrsf7* allele (Fig. 1b). Immunofluorescence analysis of spleen cells showed expression of CD27 in WT mice on the great majority of both CD4⁺ and CD8⁺ T cell subsets and a small proportion of B cells, as reported previously²⁰. It also confirmed the CD27-deficient phenotype (Fig. 1c). CD27^{-/-} mice proved viable, fertile and remained healthy over prolonged periods of time (>6 months).

T cell division upon TCR stimulation

Mice were backcrossed seven or more times to C57BL/6 and 129/Ola backgrounds. In all experiments, CD27^{+/+} and CD27^{-/-} littermates were compared. CD27^{+/+}, CD27^{+/-} and CD27^{-/-} mice did not differ in bone marrow, thymus, spleen or lymph node cellularity. Also, the T or B cell, and CD4⁺ or CD8⁺ subset, composition of




Figure 2. CD27 does not affect T cell division. T cells were purified from lymph nodes of WT and CD27^{-/-} littermates back crossed to a 129/OLA background. (a) They were cultured for 3 days in the presence of the indicated concentrations of coated mAb to CD3 ϵ . In selected wells, mAb to CD28 or recombinant IL-2 were added. [³H]thymidine incorporation proceeded for 8 h at

the end of the culture period. (Values are the mean \pm s.d. of triplicate cultures in one experiment. The experiment was repeated four times with similar results.) (b) Purified T cells were labeled with the fluorescent dye CFSE before incubation in the presence of coated mAb to CD3 ϵ and mAb to CD28, or CD27 as indicated. After 3 days, cells were collected and analyzed by FACS after gating of live cells.

not influence cell division. The CFSE-staining pattern obtained was identical to the pattern obtained after stimulation with mAb to CD3 alone (Fig. 2b). Lack of effect of mAb to CD27 cannot be explained by lack of CD27 expression: the great majority of naive peripheral T cells express CD27 (Fig. 1c) and costimulation with mAb to CD27 enhances [³H]thymidine incorporation three- to fourfold²⁰. We conclude that, unlike CD28, CD27 does not influence the cell division process. Its effect on [³H]thymidine incorporation might be explained by an effect on cell survival.

Primary T cell response in CD27^{-/-} mice

To test the capacity of the immune system to mount an antigen-specific T cell response in the absence of CD27-CD70 interaction, we studied influenza virus infection in C57BL/6 mice. Major histocompatibility complex (MHC) H2-D^b tetramers loaded with the immunodominant influenza NP(366–374) peptide allow visualization and quantification of virus-specific CD8⁺ T cells^{33,34}. Mice were infected intranasally with 25 hemagglutinin units (HAU) of influenza virus A/NT/60/68. At days 6, 8, 10 and 14 after infection, absolute numbers of lung-infiltrating CD4⁺ and CD8⁺ T cells were measured (Fig. 3a). In WT mice, T cell numbers were at a maximum 8–10 days after infection, with CD8⁺ cells outnumbering CD4⁺ cells. In CD27^{-/-} mice, the overall response curves were lower than in WT mice, for both CD4⁺ T cells ($P=0.014$) and CD8⁺ T cells ($P=0.020$).

We determined the percentage and absolute numbers of virus-specific CD8⁺ T cells in lung and spleen using MHC class I tetramers. A representative tetramer *versus* CD8 staining of lung-infiltrating cells is shown in Fig. 3b. In the upper panels of Fig. 3c, tetramer-

positive T cells are plotted as percentage of total CD8⁺ T cells. At the peak of the response (days 8–10) in WT mice, 10–15% of CD8⁺ T cells in the lung and 4–5% of CD8⁺ T cells in the spleen were specific for the immunodominant influenza virus peptide. This translated into 20–30 $\times 10^4$ virus-specific T cells in the lung and around 6 $\times 10^5$ virus-specific T cells in the spleen. In lung, a difference in responsiveness between WT and CD27-deficient mice was not apparent from the percentage of tetramer-positive T cells, but it became evident when absolute numbers were calculated based on the size of the infiltrate (Fig. 3c, lower panels). The curve for absolute numbers of virus-specific CD27^{-/-} T cells shifted downwards, compared to the curve for CD27^{+/+} T cells, for both lung ($P=0.0046$) and

spleen ($P=0.014$). Thus, CD27^{-/-} mice displayed a decrease in absolute numbers of CD8⁺ and CD4⁺ T cells that infiltrate the lung in response to primary infection and a reduction in absolute numbers of virus-specific CD8⁺ T cells in lung and spleen, particularly at the peak of the response.

T cell memory in the absence of CD27

To analyze the memory response, CD27^{-/-} and WT mice were challenged with 200 HAU of influenza virus 6 weeks after the first infection. In the lungs of WT mice, the secondary virus-specific T cell response could be measured earlier than the primary, with the peak of the response at day 5 (Fig. 4a). Moreover, the response was more pronounced with around 40% of CD8⁺ T cells (60×10^4 cells) in the lung being virus-specific at the peak (Fig. 4b). Quantification of total CD4⁺ and CD8⁺ T cells present in the lungs at days 3, 5, 7 and 11 after reinfection revealed that the response was reduced in CD27^{-/-} mice (Fig. 4a). For CD4⁺ T cells, response kinetics in CD27^{-/-} mice were delayed ($P=0.053$) as compared to WT mice, with a peak at day 7 instead of day 5. At all time points, lung-infiltrating CD4⁺ T cell numbers in CD27^{-/-} mice were below those found in WT mice. Also for CD8⁺ T cells, the kinetics of the response were markedly delayed in CD27^{-/-} mice ($P=0.0007$), with the peak of the response at day 7 instead of at day 5. The curve was also shifted downwards ($P<0.0001$): CD8⁺ T cell numbers in lungs of CD27^{-/-} mice were below those in WT mice at days 3 and 5 and peak values remained at the level seen in the primary response (Fig. 3a).

The discrepancy in responsiveness of WT and CD27^{-/-} mice could




Figure 3. CD27 is important for generation of T cell immunity to influenza virus. CD27^{+/+} (solid lines, filled squares) and CD27^{-/-} (broken lines, open squares) littermates were infected intranasally with 25 HAU influenza A virus strain NT/60/68 and killed 6, 8, 10 or 14 days later. Three to six mice were used for each time point. Cells were isolated from lungs and spleen, counted, stained with mAb to CD4 or double stained with mAb to CD8 and recombinant H2-D^b tetramers, and analyzed by flow cytometry. (Data points are from individual mice, dashes indicate means.) (a) Absolute numbers of CD4⁺ and CD8⁺ T cells in the lungs at the indicated days after primary infection. (b) Representative FACS profile of T cells in lung stained with MHC class I tetramers and mAb to CD8. (c) MHC class I tetramer-positive T cells in lung and spleen at the indicated time points after primary infection, expressed as percentage of CD8⁺ cells and in absolute numbers.

also be seen in the virus-specific CD8⁺ T cell infiltrate of the lung (Fig. 4b). As for the primary response, the percentage of tetramer-positive T cells did not indicate a major difference in responsiveness between WT and CD27-deficient mice. However, when

absolute numbers were calculated based on the size of the lung infiltrate, a vast discrepancy became apparent. At day 5, a peak response was found in WT mice, which was lacking in CD27^{-/-} mice ($P=0.0003$). The response in CD27^{-/-} mice was comparable to the




Figure 4. CD27 is required for T cell memory. Six weeks after primary infection, CD27^{+/+} (solid lines, filled squares) and CD27^{-/-} (dashed lines, open squares) were infected intranasally with 200 HAU influenza A virus and killed 3, 5, 7 or 11 days later. (Three to six mice were used for each time point. Cells were isolated and stained as described for Fig. 3. Data points are from individual mice, dashes indicate means.) (a) Absolute numbers of CD4⁺ and CD8⁺ T cells in the lungs at the indicated days after secondary infection. (b) MHC class I tetramer-positive T cells in lung and spleen at the indicated time points after secondary infection, expressed as percentage of CD8⁺ cells and in absolute numbers.




Figure 5. CD27 contributes to the antigen-specific CD4⁺ T cell response.

Expansion of splenic CD4⁺ T cells upon challenge with purified, inactivated influenza virus *in vitro*. CD4⁺ T cells were purified and cultured for 4 days with the indicated concentrations of virus, presented by C57BL/6 spleen cells. [³H]thymidine incorporation occurred during the last 8 h of culture. (Values represent the mean \pm s.d. of triplicate samples in one experiment. The experiment is representative of three.) T cells were taken from immunized CD27^{+/+} or CD27^{-/-} mice or from nonimmunized (N) WT mice at day 14 after primary infection (upper panel) or day 11 after secondary infection (lower panel).

primary response in terms of percentage (about 15%) and absolute number (about 7×10^4) of tetramer-positive CD8⁺ T cells at the peak. In the spleen, the curve for CD27^{-/-} mice was shifted downwards compared to the curve for WT mice ($P=0.0022$). In conclusion, the memory T cell response proceeded with slower kinetics in CD27^{-/-} mice and was greatly reduced in terms of absolute numbers of participating CD8⁺ T cells, particularly in the lung.

Virus-specific CD4⁺ T cell responses

Absolute numbers of CD4⁺ T cells infiltrating the lung after primary and secondary virus infection were reduced in CD27^{-/-} mice. To examine whether the antigen-specific CD4⁺ T cell response was affected, we assayed for virus-specific CD4⁺ T cell expansion *in vitro*. Purified CD4⁺ T cells from both CD27^{+/+} and CD27^{-/-} mice showed a specific expansion, dependent on the concentration of viral antigen (Fig. 5). The response of CD27^{-/-} CD4⁺ T cells was reduced as compared to the response of WT cells, both after primary and secondary infection, over a wide concentration range of viral protein. This supported the *in vivo* response data, indicating that both primary and memory CD4⁺ T cell responses were affected in CD27^{-/-} mice.

Table 1. CD27 does not affect differentiation towards IFN- γ production

		CD8 ⁺	CD4 ⁺
Naive	+/+	8.2 \pm 1.5 ¹	11.5 \pm 5.0
	-/-	5.9 \pm 0.3	9.4 \pm 0.1
Primary	+/+	12.2 \pm 5.2	7.2 \pm 3.6
	-/-	10.8 \pm 3.2	10.3 \pm 7.5
Secondary	+/+	19.9 \pm 8.2	18.3 \pm 9.7
	-/-	20.7 \pm 11.3	18.8 \pm 13.2

Lung-infiltrating cells were stimulated with PMA and ionomycin. IFN- γ -positive cells were quantified within gated CD8⁺ or CD4⁺ cell populations. Results are percentages expressed as mean \pm s.d. derived from three independent experiments with two mice each per group.

CD8⁺ T cell differentiation into effector cells

To examine whether CD27 signaling contributed to differentiation of peripheral T cells into effector cells, the capacity of CD27^{-/-} T cells to exert cytotoxicity and to produce interferon γ (IFN- γ) were tested. For cytotoxicity assays, splenocytes from immunized mice were cultured *in vitro* with NP(366–374) peptide and IL-2 to allow the peptide-specific CD8⁺ T cell population to expand. A similar amount of virus-specific CD8⁺ T cells from WT and CD27^{-/-} mice were used. T cells from CD27^{-/-} mice lysed their targets as efficiently as WT controls, after both primary or secondary infection (see Web Figure 1 on the supplementary information page of *Nature Immunology* online). Thus, a lack of CD27 does not affect the differentiation program that shapes cytolytic T cell function.

The capacity to produce IFN- γ was studied in both CD4⁺ and CD8⁺ T cells isolated from mice immunized 8 days previously with influenza virus. Total lung-infiltrating cells were stimulated with phorbol ester and ionomycin and IFN- γ was detected by intracellular staining (Table 1). No difference in the percentage of IFN- γ -producing T cells was found in CD27^{+/+} and CD27^{-/-} mice. This indicated that a similar proportion of CD4⁺ and CD8⁺ T cells present in the lungs of WT and CD27-deficient mice had completed the differentiation program that allows IFN- γ production.

Discussion

Our results show that CD27 is not essential for generation and maintenance of T and B lymphocyte populations. Previously, we hypothesized that CD27 acts as a costimulus for pre-TCR function³⁵, based on the finding that *in vivo*-administered mAb to CD27 inhibited generation of the CD4⁺CD8⁺ thymic compartment. However, T cell development was normal in CD27-deficient mice so we inferred that CD27-CD70 interaction was not required to support pre-TCR function. In our earlier experiments³⁵, mAb to CD27 presumably induced a CD27 function that counteracted the pre-TCR-induced response.

The CFSE assay showed that lack of CD27 expression in CD27-deficient cells or stimulation of CD27 with specific mAb in WT cells does not affect cell cycle activity of activated peripheral T cells. Yet both interventions affect [³H]thymidine incorporation²⁰ (Fig. 2a). This can be explained by the notion that incorporation of [³H]thymidine is not only determined by cell cycle activity, but also by the number of cells participating in the response. The latter is determined by entry into cell cycle and by cell survival. As we have clearly demonstrated that CD27 does not affect the cell division process, reduced [³H]thymidine incorporation in CD27^{-/-} T cells is most likely due to decreased T cell survival.

In contrast to CD28, CD27 employs Traf molecules to induce downstream signals, in particular Traf-2 and Traf-5^{36,37}. Whereas in Traf-5^{-/-} mice, lymphocyte homeostasis does not seem to be affected³⁸, Traf-2^{-/-} mice are severely lymphopenic, implicating this molecule in T and B cell survival⁷. Analysis of the virus-specific response of CD4⁺ T cells, isolated from the spleens of immunized mice confirmed that CD27 is essential for adequate expansion of antigen-activated T cells.

Studies suggest that CD27 may contribute to the generation of cytolytic effector cells^{13,30}. However, in these experiments, stimulatory effects of CD27 on T cell expansion may have obscured the results as the actual amount of antigen-specific T cells entered into the cytotoxicity assay was not determined. In our experiments equal amounts of peptide-specific T cells were used, which allowed the conclusion that CD27 does not stimulate the differentiation program

per se. Similarly, we found that CD27-deficient T cells of both CD8⁺ and CD4⁺ phenotype can adequately synthesize IFN- γ .

Primary influenza virus-specific responses of both CD4⁺ and CD8⁺ T cells were affected in CD27^{-/-} mice. The absolute numbers of T cells infiltrating the lung, the number of tetramer-positive CD8⁺ T cells in lung and spleen and the anti-viral response of splenic CD4⁺ T cells *in vitro* provided evidence for this conclusion. Among Traf-linked TNF receptor family members, CD40³⁹⁻⁴¹, 4-1BB^{42,43} and OX-40^{44,45} are implicated in T cell priming. CD40 plays a role on DCs, interacting with CD40L on both CD4⁺ and CD8⁺ T cells³⁹⁻⁴¹. Apart from skewing cytokine production by DC, it serves to up-regulate the ligands for CD28, such that costimulation of the T cell can take place⁴. Yet on infection with lymphocytic choriomeningitis virus (LCMV), which causes a systemic infection, the cytotoxic and helper T cell responses are not critically dependent on either CD28⁴⁶ or CD40L^{47,48}. A role of the different costimulatory molecules may be revealed depending on the model system chosen. For instance, after localized infection with influenza virus, primary CD8⁺ T cell responses are reduced in the absence of CD28 function^{49,50}.

The molecules 4-1BB and OX-40 are found on activated T cells and can interact with ligand on DCs^{24,25}. Mice that are 4-1BBL^{-/-} can generate LCMV-specific CTLs and clear the virus^{42,43}, but virus-specific CD8⁺ T cell numbers were reduced⁴³. In OX-40^{-/-} mice, the number of lung-infiltrating CD4⁺, but not CD8⁺ T cells was reduced after influenza virus infection. Splenic CD4⁺ T cells showed deficient expansion and cytokine production in response to both influenza virus and LCMV. However, primary CTL responses to these viruses were normal⁴⁴. Also, CD4⁺ T cells taken from OX-40L-deficient mice immunized with protein antigens showed impaired responsiveness upon rechallenge *in vitro*. In this assay, OX-40L^{-/-} antigen presenting cells were nonfunctional, pinpointing the defect to T cell-DC communication⁴⁵. Other assays in OX-40L^{-/-} mice revealed a defect in CD8⁺ T cell-dependent contact hypersensitivity⁵¹. Apparently, the OX-40-OX-40L pair is involved in certain, but not all, T cell-DC interactions and is not selective for CD4⁺ or CD8⁺ T cells. CD70 has been detected on activated mouse DCs²⁷. Therefore, the CD27-CD70 pair may contribute to T cell priming at the DC-T cell interphase, like 4-1BB and OX-40. In addition, both CD70 and OX-40L are expressed on activated T cells and can support T cell expansion during T-T cell interaction.

The most dramatic defect in CD27^{-/-} mice concerned T cell memory, in particular of CD8⁺ T cells. Memory is dependent on the amount of surviving T cells after primary TCR-mediated activation and presumably on escape from activation-induced cell death. Traf-linked TNF receptor family members, by virtue of their anti-apoptotic effects, would be prime candidates to shape T cell memory. However, thus far, only CD40L has been implicated in T cell memory using gene-targeted mice⁵². OX-40 was shown to inhibit peripheral deletion and enhance long-term T cell survival by *in vivo* antibody injection studies⁵³. CD27 may exert its effects by promoting T cell survival and/or expansion of memory T cells. We cannot exclude a role for CD27 in lymphocyte trafficking. Recently, its close relative OX-40 was connected to cell migration. OX-40 was found to up-regulate expression of the CXCR5 chemokine receptor and allow T cell migration into B cell areas of the peripheral lymphoid organs⁵⁴.

Although the CD27^{-/-} mice had reduced T cell immunity, in particular the CD8⁺ T cells, they did not become overtly more ill and recovered from influenza virus infection within the same time peri-

od as WT mice. In part, this may be explained by the finding that CD27^{-/-} mice have a normal B cell response to influenza virus. Serum titers of virus-specific immunoglobulins of all isotypes are in the same order of magnitude as in WT mice (data not shown). Also, immunoglobulin levels remain high in both CD27^{-/-} mice and WT mice for more than 3 months after primary infection and therefore are expected to contribute to virus elimination after secondary challenge.

This study underlines the importance of Traf-linked TNF receptor family members in induction and maintenance of the specific immune response. These molecules provide a lead for the development of alternative, subtle immunological intervention for conditions of excessive or deficient immune responsiveness, such as autoimmunity and cancer.

Methods

Mice. All mice were bred in the facility of The Netherlands Cancer Institute under specific pathogen-free conditions and used for experiments at 6 to 8 weeks of age. All animal experiments were carried out according to institutional and national guidelines and approved by the Experimental Animal Committee of The Netherlands Cancer Institute (DEK).

Generation of CD27-deficient mice. Clone λ 5 was isolated from a genomic 129/OLA DNA library in phage λ FIX (Stratagene, La Jolla, CA) by probing with a 5' 600-bp *EcoRI-PstI* fragment of the murine CD27 cDNA¹². *XbaI* and *XbaI-NorI* fragments of λ 5 were subcloned and characterized by Southern blotting with CD27 cDNA and by nucleotide sequencing. To make the targeting construct, a λ 5 fragment containing the entire coding region, from the *BglII* site just upstream of the ATG codon to the *BamHI* site, downstream of the stop codon, was replaced by the hygromycin resistance gene under the control of the GK promoter. Subclone IB10, derived from the 129/OLA ES cell clone E14 was electroporated with a *NorI-Sall* fragment of the targeting construct. Genomic DNA of ES cell clones resistant to 150 μ g/ml hygromycin B (Calbiochem, La Jolla, CA) was diagnosed by Southern blotting, as described in the Results.

Genotypic analysis of mice. The mutated *Tnfrsf7* allele was detected by PCR on tail DNA of the hygromycin B-resistance gene, using nucleotides 256-274 as sense and 914-933 as antisense primers. The *Tnfrsf7* WT allele was detected using nucleotides 869-889 of the cDNA as sense and 1236-1255 as antisense primers¹².

Flow cytometry. Peripheral blood was collected and cells were spun down. Erythrocytes were then lysed in 0.14 M NH₄Cl, 0.017 M Tris-HCl at pH 7.2 on ice for 10 min. Peripheral blood cells and cells from bone marrow, thymus, spleen and lymph nodes were preincubated with an Fc Block (mAb to CD16-CD32, 2.4G2, PharMingen, San Diego, CA) and washed in FACS buffer (PBS, 0.5% bovine serum albumin, 0.01% sodium azide). Next, they were incubated with specific antibody directly conjugated to fluorescein isothiocyanate (FITC), phycoerythrin (PE) or biotin as indicated, washed, incubated with Tricolor-streptavidin or PE-streptavidin (Caltag, San Diego, CA) when applicable, washed and analyzed using a FACScan and LYSYS II software (Becton Dickinson, Mountain View, CA). Immunofluorescence mAbs were from PharMingen or donated by investigators. The mAbs we used were anti-CD3e, 500A2; anti-CD4, RM4-5; anti-CD8 α , 53-6.7; anti-CD27, LG.3A10³⁰; anti-CD28, 37.51; anti-CD44, IM7; anti-CD45R (B220), RA3.6B2; anti-CD69, H1.2F3; anti-IgM, R6-60.2; anti-IFN- γ , XMG1.2. PE-labeled tetramers of the murine class I molecule H2-D^b, β_2 -microglobulin and the influenza nucleoprotein peptide ASNENMDAM, NP(366-374), were prepared as described³². Cells were incubated with MHC class I tetramers together with FITC-anti-CD8 α . Cells were washed in FACS buffer and analyzed using a FACSCalibur and CELLQuest software (Becton Dickinson, Mountain View, CA).

Preparation of purified T cells. Lymph nodes and spleens were dissociated in Iscove's Modified Dulbecco's medium supplemented with 8% fetal calf serum (FCS), penicillin and streptomycin. Erythrocytes were lysed in 0.14 M NH₄Cl, 0.017 M Tris-HCl, pH 7.2 for 5 min on ice. Cells were washed in medium, passed over nylon wool (Polysciences, Warrington, MA) and incubated on ice for 30 min with mAb M5/114 to MHC class II. In case of CD4⁺ T cell purification, 2 μ g of mAb to CD8 α per 10⁶ cells was added. This was followed by a 30-min incubation on ice with 100 μ l of goat anti-mouse immunoglobulin-coated magnetic beads and 20 μ l of goat anti-rat immunoglobulin-coated magnetic beads per 10⁷ cells (Advanced Magnetics Inc., Cambridge, MA). Purity of the resulting cell population was checked by immunofluorescence analysis with mAbs to CD3e and B220. Only preparations that contained more than 98% T cells or CD4⁺ T cell preparations that were more than 90% pure were used.

***In vitro* T cell stimulation.** Purified T cells from backcrossed mice (10⁵ per well) were plated in IMDM with 8% FCS in 96-well round-bottom plates, coated with mAb 145-

2C11 to CD3ε at the indicated concentrations, in the presence or absence of coated mAb 37.51 to CD28 (10 μg/ml) or IL-2 (100 U/ml). Cells were cultured for 3 days. During the last 8 h of culture, 0.4 μCi [³H]thymidine (Amersham International, Amersham, UK) was present per well. Proliferation of virus-specific CD4⁺ T cells was determined using 2×10⁵ purified spleen CD4⁺ T cells per well, taken from mice at day 14 after primary infection or at day 11 after secondary infection. Sucrose gradient-purified influenza virus was inactivated with 2% paraformaldehyde in PBS for 20 min at room temperature followed by addition of glycine to 0.2 M and overnight dialysis in PBS. Virus was added to the culture at indicated concentrations. Irradiated (25 Gy) C57BL/6 spleen cells were used as antigen presenting cells at 4×10⁶ cells per well. Cells were cultured for 4 days; for the last 8 h in the presence of [³H]thymidine. Cells were collected with a Titertek automatic harvester (ICN, Costa Mesa, CA) and incorporation of radioactivity was measured in a β-plate counter (Pharmacia, Uppsala, Sweden). For cell cycle analysis, purified T cells were labeled with 5 μM of CFSE (Molecular Probes) before stimulation. Cells were stimulated with coated mAb to CD3ε with or without coated mAb to CD28 (both at 10 μg/ml), or soluble mAb to CD27 (0.1 μg/ml). Gated live cells were analyzed by flow cytometry after 3 days of culture.

Cytotoxicity assay. Cytolytic activity of virus-specific CD8⁺ T cells derived from spleens from infected mice was determined in a 5-h ⁵¹Cr-release assay. Spleens from mice at day 14 after primary infection or day 11 after secondary infection were dissociated into single cell suspensions and restimulated for 7 days in the presence of 0.5 ng/ml of NP-derived peptide ASNENMDAM and 20 U/ml of recombinant human IL-2. EL-4 target cells were preincubated with peptide for 1 h at 37 °C. Percent specific lysis was calculated from the equation:

$$\frac{(\text{experimental } ^{51}\text{Cr-release} - \text{spontaneous } ^{51}\text{Cr-release})}{(\text{maximum } ^{51}\text{Cr-release} - \text{spontaneous } ^{51}\text{Cr-release})} \times 100 = \% \text{ specific lysis.}$$

Intracellular cytokine staining. Lung-infiltrating cells were stimulated for 5 h with 5 ng/ml of phorbol 12-myristate 13-acetate and 1 μM of ionomycin in the presence of 10 μg/ml of brefeldin A (Sigma, St. Louis, MO). Cells were collected and washed once with FACS buffer. After incubation with Fc Block and staining with PE-conjugated mAb to CD4 and FITC-conjugated mAb to CD8, cells were fixed with 4% paraformaldehyde in PBS for 5 min. They were washed once with FACS buffer, containing 0.1% saponin (Sigma) for permeabilization, and incubated in the same buffer with allophycocyanin-conjugated mAb to IFN-γ for 30 min on ice. Cells were washed three times, resuspended in FACS buffer and analyzed by flow cytometry.

Virus infection. Influenza virus strain A/NT/60/68 was grown, purified and tested for hemagglutinin activity and infectious titers in the Department of Virology, Erasmus University Rotterdam, The Netherlands. Mice used in these experiments were backcrossed at least seven times to the C57BL/6 background and CD27-deficient and WT littermates were compared in all experiments. Mice were anesthetized and infected intranasally with 50 μl of PBS with or without 25 or 200 HAU of virus for primary and secondary infections, respectively. Mice were killed at indicated time points after infection, and lungs and spleen were removed. They were minced in single chamber mesh filters and cells were counted and analyzed by flow cytometry.

Statistical analysis. All statistical analyses were based on ANOVA. A one-quarter power transformation was applied to achieve normally distributed data. *P* values were calculated from type II sum of squares. Bonferroni correction (multiplication of the *P* values by 4) was used to correct for the probability of false positive findings due to use of four response measures.

Acknowledgements

We thank E. Tanger, M. Timpico, L. Tolkamp, T. Schrauwers, H. Starrevelt and other staff of the Animal Facility of the Netherlands Cancer Institute for biotechnical assistance and maintenance of the mice; A. A. M. Hart for expert statistical analysis; G. Rimmelzwaan for virus preparations and advice; E. Noteboom and A. Pfautz for assistance with flow cytometry; P. Krimpenfort, J. Haanen, G. Dingjan, R. Hendriks and J. Kirberg for experimental advice and assistance; and A. M. Kruisbeek for reading the manuscript. Supported by The Netherlands Organization for Scientific Research (NWO).

Received 18 August 2000; accepted 6 October 2000.

- Arch, R. H., Gedrich, R. W. & Thompson, C. B. Tumor necrosis factor receptor-associated factors (TRAFs)- a family of adapter proteins that regulates life and death. *Genes Dev.* **12**, 2821-2830 (1998).
- Gravestain, L. A. & Borst, J. Tumor necrosis factor receptor family members in the immune response. *Sem. Immunol.* **10**, 423-434 (1998).
- Anderson, D. M. *et al.* A homologue of the TNF receptor and its ligand enhance T-cell growth and dendritic-cell function. *Nature* **390**, 175-179 (1997).
- Koch, F. *et al.* High level IL-12 production by murine dendritic cells: upregulation by MHC class II and CD40 molecules and downregulation by IL-10. *J. Exp. Med.* **184**, 741-746 (1996).
- Grell, M. *et al.* Induction of cell death by tumor necrosis factor (TNF) receptor 2, CD40 and CD30: a role for TNF-R1 activation by endogenous membrane-anchored TNF. *EMBO J.* **18**, 3034-3043 (1999).
- Xu, Y., Cheng, G. & Baltimore, D. Targeted disruption of TRAF3 leads to postnatal lethality and

defective T-dependent immune responses. *Immunity* **5**, 407-415 (1996).

- Yeh, W. -C. *et al.* Early lethality, functional NF-κB activation & increased sensitivity to TNF-induced cell death in TRAF2-deficient mice. *Immunity* **7**, 715-725 (1997).
- Lee, S. Y., Reichlin, A., Santana, A., Sokol, K. A., Nussenzweig, M. C. & Choi, Y. TRAF2 is essential for JNK but not NF-κB activation and regulates lymphocyte proliferation and survival. *Immunity* **7**, 703-713 (1997).
- Rothe, M., Pan, M. G., Henzel, W. J., Ayres, T. M. & Goeddel, D. V. The TNFR2-TRAF signaling complex contains two novel proteins related to baculoviral inhibitor of apoptosis proteins. *Cell* **83**, 1243-1252 (1995).
- Liu, Z. -G., Hsu, H., Goeddel, D. V. & Karin, M. Dissection of TNF receptor 1 effector functions: JNK activation is not linked to apoptosis while NF-κB activation prevents cell death. *Cell* **87**, 565-576 (1996).
- Camerini, D., Walz, G., Loenen, W. A. M., Borst, J. & Seed, B. The T cell activation antigen CD27 is a member of the NGF/TNF receptor gene family. *J. Immunol.* **147**, 3165-3169 (1991).
- Gravestain, L. A. *et al.* Cloning and expression of murine CD27: comparison with 4-1BB another lymphocyte-specific member of the nerve growth factor receptor family. *Eur. J. Immunol.* **23**, 943-950 (1993).
- Goodwin, R. G. *et al.* Molecular and biological characterization of a ligand for CD27 defines a new family of cytokines with homology to tumor necrosis factor. *Cell* **73**, 447-456 (1993).
- Bowman, M. R. *et al.* The cloning of CD70 and its identification as the ligand for CD27. *J. Immunol.* **152**, 1756-1761 (1994).
- Tesselaar, K., Gravestain, L. A., van Schijndel, G. M. W., Borst, J. & van Lier, R. A. W. Characterization of murine CD70, the ligand of the TNF receptor family member CD27. *J. Immunol.* **159**, 4959-4965 (1997).
- Oshima, H. *et al.* Characterization of murine CD70 by molecular cloning and mAb. *Int. Immunol.* **10**, 517-526 (1998).
- Lens, S. M., Tesselaar, K., van Oers, M. H. J. & van Lier, R. A. W. Control of lymphocyte function through CD27-CD70 interactions. *Sem. Immunol.* **10**, 491-499 (1998).
- van Lier, R. A. W. *et al.* Tissue distribution and biochemical and functional properties of Tp55 (CD27), an novel T cell differentiation antigen. *J. Immunol.* **139**, 1589-1596 (1987).
- Hintzen, R. Q., de Jong, R., Lens, S. M. A. & van Lier, R. A. W. CD27: marker and mediator of T-cell activation. *Immunol. Today* **15**, 307-311 (1994).
- Gravestain, L. A., Nieland, J. D., Kruisbeek, A. M. & Borst, J. Novel mAbs reveal potent co-stimulatory activity of murine CD27. *Int. Immunol.* **7**, 551-557 (1995).
- Hamann, D. *et al.* Phenotypic and functional separation of memory and effector human CD8⁺ T cells. *J. Exp. Med.* **186**, 1-12 (1997).
- Maurer, D. *et al.* IgM and IgG but not cytokine secretion is restricted to the CD27⁺ B lymphocyte subset. *J. Immunol.* **148**, 3700-3705 (1992).
- Klein, U., Rajewsky, K. & Kuppers, R. Human immunoglobulin (IgM) IgD⁺ peripheral blood B cells expressing the CD27 cell surface antigen carry somatically mutated variable region genes: CD27 as a general marker for somatically mutated (memory) B cells. *J. Exp. Med.* **188**, 1679-1689 (1998).
- Vinay, D. S. & Kwon, B. S. Role of 4-1BB in immune responses. *Sem. Immunol.* **10**, 481-489 (1998).
- Weinberg, A. D., Vella, A. T. & Croft, M. OX-40: life beyond the effector T cell stage. *Sem. Immunol.* **10**, 471-480 (1998).
- Hintzen, R. Q. *et al.* CD70 represents the human ligand for CD27. *Int. Immunol.* **6**, 477-480 (1994).
- Akiba, H. *et al.* Critical contribution of OX40 ligand to T helper cell type 2 differentiation in experimental leishmaniasis. *J. Exp. Med.* **191**, 375-380 (2000).
- Kobata, T., Agematsu, K., Kameoka, J., Schlossman, S. F., Morimoto, C. CD27 is a signal-transducing molecule involved in CD45RA⁺ naive T cell costimulation. *J. Immunol.* **153**, 5422-5432 (1994).
- Hintzen, R. Q. *et al.* Engagement of CD27 with its ligand CD70 provides a second signal for T cell activation. *J. Immunol.* **154**, 2612-2623 (1995).
- Brown, G. R., Meek, K., Nishioka, Y. & Thiele, D. L. CD27-CD27ligand/CD70 interactions enhance alloantigen-induced proliferation and cytolytic activity in CD8⁺ T lymphocytes. *J. Immunol.* **154**, 3686-3695 (1995).
- Agematsu, K. *et al.* Generation of plasma cells from peripheral blood memory B cells: Synergistic effect of interleukin-10 and CD27/CD70 interaction. *Blood* **91**, 173-180 (1998).
- Lenschow, D. J., Walunas, T. L. & Bluestone, J. A. CD28/B7 system of T cell co-stimulation. *Annu. Rev. Immunol.* **14**: 233-258 (1996).
- Flynn, K. J. *et al.* Virus-specific CD8⁺ T cells in primary and secondary influenza pneumonia. *Immunity* **8**, 683-691 (1998).
- Haanen, J. B. A. G., Wolkers, M., Kruisbeek, A. M. & Schumacher, T. N. M. Selective expansion of cross-reactive CD8⁺ memory T cells by viral variants. *J. Exp. Med.* **190**, 1319-1328 (1999).
- Gravestain, L. A., van Ewijk, W., Ossendorp, F. & Borst, J. CD27 cooperates with the pre-T cell receptor in the regulation of murine T cell development. *J. Exp. Med.* **184**, 675-685 (1996).
- Gravestain, L. A. *et al.* The Tumor Necrosis Factor receptor family member CD27 signals to Jun N-terminal kinase via Traf-2. *Eur. J. Immunol.* **28**, 2208-2216 (1998).
- Akiba, H. *et al.* CD27, a member of the Tumor Necrosis Factor receptor superfamily, activates NF-κB and Stress-activated protein kinase/c-Jun N-terminal kinase via TRAF-2, TRAF-5 and NF-κB-inducing kinase. *J. Biol. Chem.* **273**, 13353-13358 (1998).
- Nakano, H. *et al.* Targeted disruption of Traf5 gene causes defects in CD40- and CD27-mediated lymphocyte activation. *Proc. Natl. Acad. Sci. USA* **96**, 9803-9808 (1999).
- Grewal, I. S., Xu, J. & Flavell, R. A. Impairment of antigen-specific T cell priming in mice lacking CD40 ligand. *Nature* **378**, 617-620 (1995).
- Schoenberger, S. P., Toes, R. E. M., van der Voort, E. I. H., Offringa, R. & Melief, C. J. M. T-cell help for cytotoxic T lymphocytes is mediated by CD40-CD40L interactions. *Nature* **393**, 480-483 (1998).
- Bennett, S. R. M. *et al.* Help for cytotoxic-T-cell responses is mediated by CD40 signalling. *Nature* **393**, 478-480 (1998).
- DeBenedette, M., Wren, T., Bachmann, M., Ohashi, P. S., Barber, B. H., Stocking, K. L., Peschon, J. J. & Watts, T. Analysis of 4-1BB Ligand (4-1BBL)-deficient mice and of mice lacking both 4-1BBL and CD28 reveals a role for 4-1BBL in skin allograft rejection and in the cytotoxic T cell response to influenza virus. *J. Immunol.* **163**, 4833-4841 (1999).
- Tan, J. T., Whitmire, J. K., Ahmed, R., Pearson, T. C. & Larsen, C. P. 4-1BB ligand, a member of the TNF family, is important for the generation of antiviral CD8 T cell responses. *J. Immunol.* **163**, 4859-4868 (1999).

44. Kopf, M., Ruedl, C., Schmitz, N., Gallimore, A., Lefrang, K., Ecabert, B., Odermatt, B. & Bachmann, M. F. OX40-deficient mice are defective in Th cell proliferation but are competent in generating B cell and CTL responses after virus infection. *Immunity* **11**, 699–708 (1999).
45. Murata, K., Ishii, N., Takano, H., Miura, S., Ndhlovu, L. C., Nose, M., Noda, T. & Sugamura, K. Impairment of antigen-presenting cell function in mice lacking expression of OX40 ligand. *J. Exp. Med.* **191**, 365–374 (2000).
46. Shahinian, A. *et al.* Differential T cell costimulatory requirements in CD28-deficient mice. *Science* **261**, 609–612 (1993).
47. Whitmire, J. K. *et al.* CD40 ligand-deficient mice generate a normal primary cytotoxic T-lymphocyte response but a defective humoral response to a viral infection. *J. Virol.* **70**, 8375–8381 (1996).
48. Oxenius, A. *et al.* CD40-CD40 ligand interactions are critical in T-B cooperation but not for other anti-viral CD4⁺ T cell functions. *J. Exp. Med.* **183**, 2209–2218 (1996).
49. Liu, Y., Wenger, R. H., Zhao, M. & Nielsen, P. J. Distinct costimulatory molecules are required for the induction of effector and memory cytotoxic T cell responses. *J. Exp. Med.* **185**, 251–262 (1997).
50. Lumsden, J. M., Roberts, J. M., Harris, N. L., Peach, R. J. & Ronchese, F. Differential requirement for CD80 and CD80/CD86-dependent costimulation in the lung immune response to influenza virus. *J. Immunol.* **164**, 79–85 (2000).
51. Chen, A. I., McAdam, A. J., Buhmann, J. E., Scott, S., Lupher, M. L., Greenfield, E. A., Baum, P. R., Fanslow, W. C., Calderhead, D. M., Freeman, G. J. & Sharpe, A. H. Ox40-ligand has a critical costimulatory role in dendritic cell:T cell interactions. *Immunity* **11**, 689–698 (1999).
52. Borrow, P., Tishon, A., Lee, S., Xu, J., Grewal, I. S., Oldstone, B. A. & Flavell, R. A. CD40L-deficient mice show deficits in antiviral immunity and have an impaired memory CD8⁺ CTL response. *J. Exp. Med.* **183**, 2129–2142 (1996).
53. Maxwell, J. R., Weinberg, A., Prell, R. A. & Vella, A. T. Danger and OX40 receptor synergize to enhance memory T cell survival by inhibiting peripheral deletion. *J. Immunol.* **164**, 107–112 (2000).
54. Walker, L. S. K. *et al.* Compromised OX40 function in CD28-deficient mice is linked with failure to develop CXC chemokine receptor 5-positive CD4 cells and germinal centers. *J. Exp. Med.* **190**, 1115–1122 (1999).